



**University of
Zurich^{UZH}**

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2019

Molecular Links Between Alzheimer's Disease and Gastrointestinal Microbiota: Emphasis on *Helicobacter pylori* Infection Involvement

Katsinelos, Taxiarchis ; Doulberis, Michael ; Polyzos, Stergios A ; Papaefthymiou, Apostolis ;
Katsinelos, Panagiotis ; Kountouras, Jannis

Abstract: Alzheimer's disease (AD) is a neurodegenerative disease and the main form of dementia, characterized by progressive cognitive decline and detrimental consequences in both personal-family and global level. Within this narrative review, we provide recent molecular aspects of Tau, a microtubule AD-associated protein, as well as amyloid beta, involved in AD pathophysiology. Moreover, we provide additional emerging data from basic research as well as clinical studies indicating an implicating role of gastrointestinal microbiota (GI-M), including *Helicobacter pylori* infection (Hp-I), in AD pathophysiology. Likewise, we identified through a molecular prism the current evidence of AD pathogenesis as well as its linkage with GI-M and emphasizing the role of Hp-I. All in all, additional large-scale studies are required for the further clarification of AD pathophysiology and its connection with GI-M and Hp-I, so as novel therapies on molecular basis become available.

DOI: <https://doi.org/10.2174/1566524019666190917125917>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-181211>

Journal Article

Accepted Version

Originally published at:

Katsinelos, Taxiarchis; Doulberis, Michael; Polyzos, Stergios A; Papaefthymiou, Apostolis; Katsinelos, Panagiotis; Kountouras, Jannis (2019). Molecular Links Between Alzheimer's Disease and Gastrointestinal Microbiota: Emphasis on *Helicobacter pylori* Infection Involvement. *Current Molecular Medicine*, 20(1):3-12.

DOI: <https://doi.org/10.2174/1566524019666190917125917>

Molecular links between Alzheimer's disease and Gastrointestinal microbiota: emphasis on *Helicobacter pylori* infection involvement

Taxiarchis Katsinelos, PhD¹, Michael Doulberis, MD, DVM, PhD^{2,3}, Stergios A Polyzos MD, MSC, to the liver and then excretion into the bile PhD^{3,4}, Apostolis Papaefthymiou MD^{3,5}, Panagiotis Katsinelos, MD, PhD³, Jannis Kountouras, MD, PhD^{3✉}

¹ UK Dementia Research Institute at the University of Cambridge, Hills Road, Cambridge, CB2 0AH, United Kingdom

^{2,3} Department of Gastroenterology and Hepatology, University of Zurich, Zurich 8091, Switzerland

³ Department of Internal Medicine, Second Medical Clinic, Ippokration Hospital, Aristotle University of Thessaloniki, Thessaloniki 54642, Macedonia, Greece

⁴ First Department of Pharmacology Medical School, Aristotle University of Thessaloniki, Thessaloniki 54124, Macedonia, Greece

⁵ Department of Gastroenterology, 401 General Military Hospital of Athens, Athens 11525, Greece

✉Address correspondence to Jannis Kountouras MD, PhD, Professor of Medicine, Gastroenterologist, Department of Medicine, Second Medical Clinic, Aristotle University of Thessaloniki, Ippokration Hospital, 8 Fanariou St, Byzantio, 551 33, Thessaloniki, Macedonia, Greece; Tel/Fax:+30-2310-892238, +30-2310-892744; E-mail: jannis@auth.gr, ancoratus2010@gmail.com

Keywords: Alzheimer's disease, tauopathies, amyloid beta, gastrointestinal microbiota, *Helicobacter pylori*

Abstract

Alzheimer's disease (AD) is a neurodegenerative disease and main form of dementia, characterized by progressive cognitive decline and detrimental consequences in both personal-family and global level. Within this narrative review, we provide recent molecular aspects of Tau, a microtubule AD-associated protein as well as amyloid beta, involved in AD pathophysiology. Moreover, we provide additional emerging data from basic research as well as clinical studies indicating an implicating role of gastrointestinal microbiota (GI-M), including *Helicobacter pylori* infection (*Hp-I*), in AD pathophysiology. Likewise, we identify through a molecular prism the current evidence of AD pathogenesis as well as its linkage with GI-M and emphasizing the role of *Hp-I*. All in all, additional large-scale studies are required for the further clarification of AD pathophysiology and its connection with GI-M and *Hp-I*, so as novel therapies on molecular basis become available.

Introduction

Alzheimer's disease (AD) is the major cause of dementia characterized by the accumulation of amyloid beta ($A\beta$) protein, the hyperphosphorylation of tau protein, and neuroinflammation, thought to lead to neuronal loss and synaptic dysfunction (1). AD is estimated to affect worldwide approximately 46.8 million people over the age of 60, having a rising tendency (2–4); its incidence and prevalence rise exponentially with increasing age particularly in the western societies (5), except preventive strategies are developed (2). The predictions regarding the prevalence of the disease, led the World Health Organization (WHO) to describe AD as a “fast-growing epidemic” (6), as a result of its prediction that the number of affected people will triple by 2050 (7).

The economic consequences as well as the traumatic burden for both patients and caregivers are enormous. The outcome is fatal and patients develop with the time impaired cognition with parallel continuously deteriorated ability to carry out daily activities and onset of neuropsychiatric symptoms (8). The most characteristic symptom at the initial phase of the disease is the loss of short-term memory (9). During disease progression, the clinical manifestations are further expanded and include speaking problems, sudden changes in mood, depression and severe orientation problems (9,10). The later stages of AD are characterized by complete withdraw from family and social activities and finally, loss of body functions that ultimately lead to death. Importantly, the constantly increasing life expectancy due to major improvements in life quality and successful treatment of previously incurable diseases, represent a critical challenge for ageing-related syndromes (11). Especially for AD, there is a massive psychological as well as the mentioned economic burden for both patients and their families, which is reflected to the society (12).

Therefore, the utmost goal of the research community is to identify effective therapeutic strategies that do not only provide symptomatic relief (13), but also to intervene in the molecular mechanisms involved in the pathophysiology of this disease, that is complex and not yet fully elucidated. In the present narrative review, we aimed to provided new molecular

links between AD, gastrointestinal microbiota and *Helicobacter pylori* infection involved in the disease pathophysiology.

Alzheimer's disease and molecular data

AD typical histopathological findings include the extracellular senile plaque formation of amyloid- β peptide ($A\beta$) and the intracellularly accumulating Neurofibrillary Tangles (NFTs) of hyperphosphorylated tau (14).

Generally, regarding $A\beta$ deposition involved in senile plaques formation (15), the proteolytic cleavage of the Amyloid Precursor Protein (APP) by β - and γ -secretases leads to the formation of this small peptide (16,17), with its length ranging from 38- up to 43-amino acids. Even though the vast majority of the produced peptides have very fast turnover and are quickly degraded, the 42-amino acid long version ($A\beta_{42}$) is highly aggregation-prone and exhibits enhanced toxicity (18). Inside the amyloid plaques, $A\beta$ is aggregated and thereby forming not only large fibrils, but also smaller insoluble oligomeric formations (19). Importantly, these amyloid plaques are usually surrounded by degenerating neurons and an increased number of activated microglia, thereby indicating an inflammatory response (20).

Likewise, regarding the implication of tau protein in AD etiology and progression, the intracellular deposition of hyperphosphorylated tau aggregates, leads to the formation of NFTs and Paired Helical Filaments (PHFs) (21,22).

Tau-related molecular data

Tau is a microtubule (MT)-associated protein involved in both the health and disease of neurons. In humans, the Microtubule Associated Protein tau gene (MAP) is transcribed predominantly throughout the Central Nervous System (CNS) and produces a long pre-mRNA that contains 16 exons (23). Specifically, six isoforms of tau are expressed predomi-

nantly in the CNS due to differences regarding the number of the 29-amino acid residue near the aminoterminal region (N-terminal projection domain) as well as in the presence of 3- or 4-repeat domains close to the carboxy-terminal insert (MT Binding Repeat Domain, RD) (24). Being the major functional role, the interaction of tau with the MTs is mediated through the electrostatic interactions between the positively charged RD and the negatively charged MT surface (25). The reduction of the tau-MT affinity is achieved through phosphorylation of tau, which under healthy conditions is a tightly regulated process (24,26).

The major hypothesis linking AD to the observed hyperphosphorylation in AD patients is the loss of this tight regulation for tau phosphorylation, which leads the protein to a higher phosphorylation state and consequently detaches from MTs and starts aggregating (27). This is mostly based on the observation that in PHFs from AD patients tau is found to be hyperphosphorylated in epitopes that are not present in healthy subjects (28–30). Even though tau can also be found to be hyperphosphorylated in conditions not related to AD (31), it is widely accepted that abnormal phosphorylation is a major (but not the exclusive) factor contributing to tau aggregation.

Patient-derived as well as *in vitro*-assembled tau filaments demonstrate the typical for amyloids cross- β structure (32,33), with the core consisting of about 90 amino acids within the RD region. As this part of the protein is also responsible for binding to MTs, tau aggregation and MT-interaction are considered as two mutually exclusive processes (34,35). Additionally, while the RD is the core of tau filaments, the N- and C-terminal parts are forming the characteristic “fuzzy coat” that surrounds the aggregation core (36,37).

The exact mechanism of tau-induced neurotoxicity remains unclear. A reasonable hypothesis is the loss of tau function due to hyperphosphorylation and aggregation. However, tau knockout mice do not demonstrate significantly overt phenotypes (38), most probably because other MAPs compensate the loss of tau function. Another proposed mechanism is the mislocalization of tau into post-synaptic spines and thereby leading to synaptic dysfunction (39). However, the most accepted scenario for the tau-mediated toxicity observed in AD is the gain of toxic function for the aggregated species. Even though initially the NFTs were

suggested to be the detrimental factor for the pathological phenotypes in neurons, there is growing evidence towards the significance of smaller oligomeric tau species. This is mostly based on the establishment of several experimental tau models that develop toxic phenotypes due to the presence of aggregates, but in the absence of fibrillar tau (40). However, the determination of the exact species responsible for the tau-induced toxic effects is extremely challenging, mostly due to the great complexity in preparation and assay read-out heterogeneity among several experimental efforts.

Additionally, the sequential deposition of aggregated tau species throughout the brain, combined with the elevated levels of tau in the cerebrospinal fluid (CSF) before AD symptoms occur (41), the affected brain areas, and the symptoms, led to the “tau spreading” hypothesis. According to this model, misfolded tau trans-cellularly spreads from initially affected brain areas even to long distant neuronal clusters (42), thereby propagating the disease-related pathology. This process has been recapitulated in murine systems through injection of patient-derived material or *in vitro* generated tau fibrils (43–46).

Specifically, the mentioned increased detection of tau in the CSF in various forms (full-length, differentially phosphorylated, cleaved etc.) is a well-established observation in AD (47). Since CSF-tau is also found in non-demented patients with acute brain injury or stroke (48), release of tau from cells was initially empirically interpreted as a result of cell death. However, the levels of phosphorylated or non-phosphorylated tau in the CSF of AD patients are reproducibly higher compared to adjusted controls with intact cognition (49–52) and the detection precedes the occurrence of the symptoms (41). Moreover, over the last years there has been a vast number of scientific studies regarding the active release of tau from cells (53) and it is firmly established that this occurs even in healthy neurons (54). These observations, combined with the “tau spreading” hypothesis as a determining factor for disease progression, eventually raised questions regarding a potentially active mechanism for tau secretion. As tau does not contain a signal peptide sequence, its secretion seems to follow other pathways, collectively referred to as “Unconventional Protein Secre-

tion" (UPS) (55,56) and more specifically the proposed pathways for tau are divided either in Type I or Type III UPS-dependent mechanisms (57–59).

A β - related data: infection hypothesis

Regarding A β and AD, although the implication in pathogenesis is well established, a previously thought causative relationship has been questioned; A β brain depletion therapies have mostly failed, thereby signifying further efforts to investigate novel mechanisms in order to clarify the pathophysiology of the disease and introduce new interventional and/or preventive strategies. In this respect, the "infection hypothesis" was recommended, though received little attention. Nevertheless, recent data indicate that A β is an antimicrobial peptide with broad-spectrum antimicrobial properties acting against bacteria, fungi, and viruses (60). Indeed, relative data indicate that microbial infections increase the synthesis of A β and the latter appears to be an innate immune defense antimicrobial peptide, which utilizes fibrillation to protect the host from variable pathogens. Moreover, brain infections often result in augmented amyloidogenic processing of the amyloid- β protein precursor and subsequent fibrillary aggregates of A β (61). The A β oligomers exhibit strong and broad-spectrum antimicrobial actions by inducing fibrils, which entrap infectious pathogens and disrupt cell membranes (62), while A β overexpression in experimental models leads to increased resistance to infection from viruses and bacteria (63,64). Therefore, the antimicrobial role of A β may explain why increased rates of infection have been observed in some of the AD clinical trials that depleted A β . Even though the A β production might be beneficial on the initial microbial challenge, potentially has gradually detrimental consequences as the infections turn to chronic and the host endeavors to remove the excess A β may reduce over time as a result of both microbial biofilm formation and microglial senescence (60,65). These data provide evidence to an infection hypothesis involved in AD pathophysiology.

B. Gastrointestinal microbiota including *Helicobacter pylori* and AD: molecular data

The gastrointestinal microbiota (GI-M) including *Helicobacter pylori* (*Hp*) and brain relationship is assumed as a potential pathogenic basis for neurological disorders of pronounced health impact such as AD (66,67); the GI-M-brain axis is critical for maintaining GI homeostasis and its dysregulation is involved in various disease states such as AD. Specifically, the GI-M is considered the principal reservoir of microbes in the human body, containing trillions of microbes and comprises the highest one among all microbial ecosystems, as it encodes approximately 4×10^6 genes, which represents 150 times more the amount of human genes (68). The human GI-M as a vital endocrine “organ” is involved in important functions, such as influencing metabolism, inducing trophic and protective effects and instructing innate immunity (69). Moreover, several data highlight the role of the GI-M in the development of neurodegenerative diseases, and in particular AD; the contribution of GI-M dysbiosis (70) to the pathogenesis of AD is well depicted in animal models of AD (71,72). In this regard, there is a great deal of impetus for the complete understanding of the pathological function, genetic evidence and functional diversity of the GI-M that could result in AD development and the detection of relative risk factors (73).

Growing clinical evidence (74) discloses that the pathophysiology of several neurodegenerative disorders could depend on the GI-M and that the resident commensal microbiota influences neuroinflammation (75). In this regard, in a recent study, fecal samples from 43 AD patients and equal number of cognitive intact controls were collected. By using 16S ribosomal RNA sequencing technique, fecal microbiota composition was analyzed and was found to be different between the two groups; bacteria taxa such as *Bacteroides*, *Actinobacteria*, *Ruminococcus*, *Lachnospiraceae*, and *Selenomonadales* were different in AD patients from those in controls at taxonomic levels (76). Similar results revealed a study from Japan. Fecal samples from demented and non-demented patients were obtained, and the gut microbiome was assessed by terminal restriction fragment length polymorphism analysis. The

analysis disclosed differences in the composition of the gut microbiome; the number of *Bacteroides* (enterotype I) was lower and the number of 'other' bacteria (enterotype III) was higher in demented patients. Multivariable analyses demonstrated that the populations of enterotype I and enterotype III bacteria were both strongly associated with dementia, independent of the traditional dementia biomarkers (77). Other authors reported that gut microbiota (GM) of AD patients revealed decreased microbial diversity and is compositionally distinct from control participants. Moreover, they found phylum- through genus-wide differences in bacterial abundance such as decreased *Firmicutes*, increased *Bacteroidetes*, and decreased *Bifidobacterium* in the fecal microbiome of AD patients. Interestingly, the authors also observed correlations between levels of differentially abundant genera and CSF biomarkers of AD like A β 42/A β 40, phosphorylated tau (and their ratio) and chitinase-3-like protein 1 (78). Additional study demonstrated that the amount of *Eubacterium rectale* and *Bacteroides fragilis* microorganisms in patients with amyloidosis were significantly reduced compared to the cognitive intact group, whereas the number of *Escherichia/Shigella* bacteria significantly increased, thereby supporting the notion that AD affects the GM (79). Furthermore, a large multicenter study revealed, that AD patients were characterized by significantly lower concentrations of cholic acid (primary bile acid) and increased levels of the bacterially produced deoxycholic acid (secondary bile acid) as well as of its conjugated forms. The increased deoxycholic to cholic acid ratio (reflection of 7 α -dehydroxylation by GM), was strongly associated with cognitive decline (80). Likewise, rising evidence from epidemiologic studies and animal models links periodontitis, caused by oral microbiota with AD (81–83).

Focus on *Hp* infection (*Hp*-I), a study from France, assessed both clinical and biological data of 53 patients with AD, testing the suggestion that *Hp* could alter the cognitive status by increasing inflammation. Results showed a lower mini-mental state examination score ($p=0.017$), higher plasma interleukin (IL)-1 β concentrations ($p=0.025$) and increased gastric atrophy ($p=0.020$) in infected participants compared to uninfected a higher cognitive impairment in *Hp*-positive AD patients (84). In cross-sectional study, by analyzing data from the US national health and nutrition examination survey, a group from US authors also investigated

the correlation between *Hp* positivity and cognitive performance among US adults. Results showed a poorer performance of *Hp*-positive 60-90 years old participants, regarding verbal memory test compared to negative. In addition, 20-59 years old infected non-Hispanic black and women performed worse on serial digits learning total errors compared to uninfected. This was a strong demonstration that *Hp* could, in some way, correlate with cerebral function even if its causative role is still to be established (85). Other authors, proposed that *Hp* could affect AD by interacting with Galectin-3, a glycan-binding protein implicated in several physiological and pathologic processes, including cell signaling, proliferation, and migration as well as in the stimulation of immune response (86).

Concerning GI-M proteins or metabolites, they might promote neurodegeneration either by inducing amyloid development through human proteins or by increasing inflammatory reactions to endogenous neuronal amyloids (87). The exposure to bacterial amyloid proteins in the GI tract may induce priming of the immune system and promote brain autoimmunity (88,89), thereby increasing the immune response to endogenous production of neuronal amyloids involved in neurodegeneration (75).

The human innate immune system recognizes bacterial amyloid proteins by introducing many signaling pathways (90), such as the toll like receptor (TLR) 1/2, Nod-like receptor-3 protein (NLRP3), transcriptional nuclear factor kappa-B (NF- κ B), innate immune receptor CD14, and inducible nitric oxide synthase. Other misfolded proteins induced by bacteria can be involved in the tissue destruction and the production of pro-inflammatory cytokines connected with AD development (91), collectively indicating that there are interactions between brain protein misfolding and GI-M.

Disruption of the intestinal barrier (composed by mucus layer intestinal columnar epithelium and lamina propria) leads to translocation of bacteria (process known as atobiosis) and toxic substances into the bloodstream associated with critically disorders; disruption of the intestinal barrier may induce local and systemic injuries playing a role in several disorders such GI tract and liver diseases, as well as in the aging process and in the systemic in-

inflammatory response syndrome, including brain dysfunctions (92). In addition to alterations in the GM composition influencing gut permeability (93), the increased amount of bacteria in the small intestine also influences permeability, as seen in small intestinal bacterial overgrowth (SIBO) and some data indicate an increased SIBO prevalence in AD patients (94). In this respect, *Hp*-related disorders are also associated with dysmotility-induced gastrointestinal bacterial overgrowth (GIBO) that may lead to bacteremia with systemic complications including brain disorders (66). GM produces and secretes more than 100 metabolites, but their effects on the pathogenesis of AD has only begun to be elucidated. Bacterial produced acids like (iso)valeric acid, (iso)butyric, propionic, acetic and formic acid have been studied and have been shown to affect the pathogenesis of AD by disturbing the activations of microglia and astrocyte, functioning inflammatory, and against aggregations of Ab and tau (95).

Some additional data indicate that the GI-M could disrupt the integrity of the blood-brain barrier (BBB) (96,97), thereby possibly contributing to brain pathologies. Additionally, GI-M dysbiosis could increase the production of lipopolysaccharides and pro-inflammatory cytokines that combined with the activation of T lymphocytes and monocytes have been suggested to induce augmented intestinal and BBB permeability. These effects have been directly linked to subsequent misfolded proteins increase, axonal injury and neuronal demyelination involved in the pathophysiology of AD (98).

Regarding the mentioned aberrantly phosphorylated tau playing a central role in AD pathophysiology (99), recent data indicate that GI-M-induced metabolite trimethylamine N-oxide (TMAO) is increased in the CSF of patients with AD and mild cognitive impairment (MCI) (69). Moreover, CSF TMAO concentrations are related with CSF biomarkers of AD pathology (phosphorylated tau and A β) and neuronal degeneration (total tau, neurogranin, and neurofilament light chain protein), offering supplementary support for the role of GI-M in AD pathophysiology. In this respect, a healthy diet seems to reduce the risk of AD development and experimental bioactive food ingestion appears to reduce neuroinflammation, oxidative

stress, A β aggregation, and tau hyperphosphorylation with a concomitant cognition improvement (100).

GI-M and *Hp* have been shown to affect neurodegenerative processes by promoting inflammation, inducing molecular mimicry mechanisms and cerebral accumulation of A β peptides. *Hp* is one of the identified bacterial pathogens in humans, which its role is principally as a pathogen or a commensal (101). *Hp-I* has been shown to influence GI-M composition, thereby promoting dysbiosis, which represents a potential significant topic for *Hp*-related GI pathologies and AD development (102–104).

In this respect, in a mouse model mimicking human AD, transgenic APP/PS1 mice were compared to wild-type histologically, microbiologically and behaviorally. The results showed impaired spatial learning and memory appeared in APP/PS1 mice, which was further aggravated 2 months later. The brain lesions were consistent immunohistochemically with amyloid plaques. Moreover, AD histological and behavioral manifestations in the APP/PS1 mice were found to be correlated with a specific GM state and microbiota diversity of APP/PS1 mice decreased with increased age; abundance of *Helicobacteraceae* and *Desulfovibrionaceae* at the family level and *Odoribacter* and *Helicobacter* at the genus level increased significantly in APP/PS1 mice than in WT mice, while *Prevotella* abundance in WT mice was significantly higher than in APP/PS1 mice (105). Moreover, two large-scale clinical surveys demonstrate that infectious burden, among others including *Hp*, is associated with worse cognitive performance in multiethnic cohort Northern Manhattan Studies (106,107). Moreover, regarding the impact of *Hp-I* on vascular dementia, *Hp* positive patients are characterized by higher pro-inflammatory cytokine levels (tumor necrosis factor (TNF)- α , IL-1 β and IL-12) and worse scores of Montreal and Mini-Mental State Examination cognitive evaluations (108); the aforementioned pro-inflammatory cytokines appear to be involved in AD pathophysiology (109). Likewise, another recent study reported that *Hp* seropositivity is directly associated with incidence of all-cause and AD dementia and with AD mortality (110). Regarding the latter findings, by introducing the histologic analysis of gastric mucosa biopsy

for the documentation of active *Hp*-I, our own studies reported that *Hp* eradication regimen is associated with a lower 5-year mortality rate in AD patients (8). Comparable data are also obtained for glaucoma (defined as “ocular” AD) that shares similar risk factors and pathogenic mechanisms with AD (111,112). Besides, two recent published review articles summarized additional substantial studies linking AD with *Hp* and GI-M (2,102). Three main mechanisms include the possibility of *Hp* to enter in the brain via the oral-, nasal-, olfactory-pathways or through the aforementioned disrupted BBB, thereby leading to neurodegeneration; a fourth mechanism includes the possibility of *Hp*-related toxic agents to access the brain via the GI tract fast retrograde neural pathway and thus inducing brain neurodegeneration pathologies (2). Other pathogens, which have been proven isolated in AD human brain, mostly in post mortem studies, include *Chlamydophila pneumoniae*, *Borrelia burgdorferi*, and other spirochetes or *herpes simplex virus type 1* (94).

Regarding the BBB dysfunction, inflammatory mediators including cytokines and chemokines produced by *Hp*-I, induce BBB disruption. For instance, *Hp*-I releases the mentioned TNF- α , acting at a distance and involving in BBB disruption via matrix metalloproteinases upregulation (2). Likewise, *Hp*-induced Vacuolating cytotoxin (VacA) displays chemotactic activities to the bone marrow-derived mast cells (BMDMCs) and stimulates BMDMCs to produce proinflammatory cytokines that damage the BBB (113). Beyond the activated mast cells releasing vascular endothelial growth factor (VEGF), IL-8, chymase, tryptase and mast cell growth factor connected to *Hp*-I, mast cells themselves release mediators including histamine, IL-8, tryptase, and VEGF, that disturb the BBB (114). Mast cell degranulation accelerates the secretion of these potent mediators, which can orchestrate neuroinflammation and affect the BBB integrity, thereby contributing to neurodegenerative processes (66).

Finally, regarding the mentioned phosphorylated tau involved in AD pathophysiology (99), *in vitro* and animal models verified that apart from A β_{42} , *Hp* induces tau hyperphosphorylation, which is directly linked to the AD-associated neurodegeneration. Specifically, incubation of *Hp* filtrated on mouse neuroblastoma N2a cells overexpressing A β PP caused an augmented production of presenilin-2 (a component of the γ -secretase enzyme complex in-

volved in the production of A β) and A β_{42} (115). In the same study, intraperitoneal injection of *Hp* filtrate in rats induced spatial learning and memory deficits, abnormal hippocampal dendritic spine maturation, and augmented presenilin-2 and A β_{42} in rat brain hippocampus and cortex. However, *Escherichia coli* filtrate injection did not influence cognitive function in rats and A β production in rats and cells (115). Moreover, *Hp* filtrate induced substantial tau hyperphosphorylation at several AD-related tau phosphorylation sites in mouse neuroblastoma N2a cells via activation of glycogen synthase kinase-3 β (116). Equally, intraperitoneal injection of *Hp* filtrate in rats resulted in noteworthy tau hyperphosphorylation in hippocampal areas of rat brain, while microglial activation and raised brain/plasma cytokine levels were not observed. These studies appear to provide evidence of the association between *Hp*-I and AD-like A β and phospho-tau pathology, indicating that *Hp* eradication may be beneficial in the tauopathy prevention.

Collectively, viewing the aforementioned data, it seems that molecular data of GI-M and *Hp*-I may contribute to the pathophysiology of AD and thus further large-scale relative studies are necessary to elucidate this field.

C. Concluding remarks

AD is a global burden with rising tendency and affects mainly the elderly population. Since the consequences are disastrous for affected families and societies, the necessity of further investigation for elucidation of pathogenicity and development of novel therapies, is of enormous significance. Mechanistically, hallmarks of disease in CNS are regarded the extracellular amyloid plaque formation and tau protein hyperphosphorylation. The “infectious hypothesis” gains ground lately, since A β is considered as antimicrobial peptide. The GI-M, including *Hp* which its role is mainly as a pathogen or a commensal, through a plethora of emerging data from basic research as well clinical studies reinforce the aforementioned hypothesis; the interplay between brain and GI-M constitutes an axis, where immune and endocrine stimuli play a central key role for shaping homeostasis or leading to dysbiosis and

neurodegeneration. All in all, a new challenging era of understanding the complex pathophysiology of AD and impact of GI-M/*Hp* on neurodegeneration has just began. The future is expected to be exciting and further large-scale studies are warranted, so as a substantiation of our knowledge is achieved and novel etiologic therapies become available.

References

1. Onojighofia Tobore T. On the Etiopathogenesis and Pathophysiology of Alzheimer's Disease: A Comprehensive Theoretical Review. J Alzheimers Dis [Internet]. 2019 Feb 8;1–21. Available from: <https://www.medra.org/servlet/aliasResolver?alias=iospress&doi=10.3233/JAD-181052>
2. Doulberis M, Kotronis G, Thomann R, Polyzos SA, Boziki M, Gialamprinou D, et al. Review: Impact of Helicobacter pylori on Alzheimer's disease: What do we know so far? Helicobacter [Internet]. 2018 Feb;23(1):1–18. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/29181894>
3. Jan AT, Azam M, Rahman S, Almigeiti AMS, Choi DH, Lee EJ, et al. Perspective Insights into Disease Progression, Diagnostics, and Therapeutic Approaches in Alzheimer's Disease: A Judicious Update. Front Aging Neurosci [Internet]. 2017;9(NOV):356. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/29163138>
4. Hu X, Wang T, Jin F. Alzheimer's disease and gut microbiota. Sci China Life Sci [Internet]. 2016 Oct;59(10):1006–23. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27566465>
5. Qiu C, Fratiglioni L. Aging without Dementia is Achievable: Current Evidence from Epidemiological Research. J Alzheimers Dis [Internet]. 2018;62(3):933–42. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/29562544>
6. WHO. First WHO ministerial conference on global action against dementia: meeting

- report. Geneva World Heal Organ [Internet]. 2015 [cited 2018 Jan 18]; Available from: http://apps.who.int/iris/bitstream/10665/179537/1/9789241509114_eng.pdf
7. International Center at NIH F. March April 2015 Global Health Matters Newsletter. [cited 2018 Jan 18]; Available from: <https://www.fic.nih.gov/News/GlobalHealthMatters/march-april-2015/Documents/fogarty-nih-global-health-matters-newsletter-march-april-2015.pdf>
 8. Kountouras J, Boziki M, Gavalas E, Zavos C, Deretzi G, Chatzigeorgiou S, et al. Five-year survival after *Helicobacter pylori* eradication in Alzheimer disease patients. *Cogn Behav Neurol* [Internet]. 2010 Sep;23(3):199–204. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/20829670>
 9. Waldemar G, Dubois B, Emre M, Georges J, McKeith IG, Rossor M, et al. Recommendations for the diagnosis and management of Alzheimer's disease and other disorders associated with dementia: EFNS guideline. *Eur J Neurol* [Internet]. 2007 Jan;14(1):e1-26. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/17222085>
 10. McKhann GM, Knopman DS, Chertkow H, Hyman BT, Jack CR, Kawas CH, et al. The diagnosis of dementia due to Alzheimer's disease: recommendations from the National Institute on Aging-Alzheimer's Association workgroups on diagnostic guidelines for Alzheimer's disease. *Alzheimers Dement* [Internet]. 2011 May [cited 2018 Jan 18];7(3):263–9. Available from: https://ac.els-cdn.com/S1552526011001014/1-s2.0-S1552526011001014-main.pdf?_tid=a207cf90-fc38-11e7-bd49-00000aab0f02&acdnat=1516270762_612db656dcfce36495ff73d2a5d85995
 11. Holtzman DM, Morris JC, Goate AM. Alzheimer's disease: the challenge of the second century. *Sci Transl Med* [Internet]. 2011 Apr 6 [cited 2018 Jan 18];3(77):77sr1. Available from: <http://stm.sciencemag.org/content/scitransmed/3/77/77sr1.full.pdf>
 12. Winblad B, Amouyel P, Andrieu S, Ballard C, Brayne C, Brodaty H, et al. Defeating Alzheimer's disease and other dementias: a priority for European science and society. *Lancet Neurol* [Internet]. 2016 Apr [cited 2018 Jan 18];15(5):455–532. Available from:

[http://www.thelancet.com/pdfs/journals/laneur/PIIS1474-4422\(16\)00062-4.pdf](http://www.thelancet.com/pdfs/journals/laneur/PIIS1474-4422(16)00062-4.pdf)

13. Schneider LS, Mangialasche F, Andreasen N, Feldman H, Giacobini E, Jones R, et al. Clinical trials and late-stage drug development for Alzheimer's disease: an appraisal from 1984 to 2014. *J Intern Med* [Internet]. 2014 Mar [cited 2018 Jan 18];275(3):251–83. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3956752/pdf/nihms558402.pdf>
14. Ittner LM, Götz J. Amyloid- β and tau--a toxic pas de deux in Alzheimer's disease. *Nat Rev Neurosci* [Internet]. 2011 Feb;12(2):65–72. Available from: <http://dx.doi.org/10.1038/nrn2967>
15. Sadigh-Eteghad S, Sabermarouf B, Majdi A, Talebi M, Farhoudi M, Mahmoudi J. Amyloid-beta: a crucial factor in Alzheimer's disease. *Med Princ Pract* [Internet]. 2015 [cited 2018 Jan 18];24(1):1–10. Available from: <https://www.karger.com/Article/Pdf/369101>
16. De Strooper B, Saftig P, Craessaerts K, Vanderstichele H, Guhde G, Annaert W, et al. Deficiency of presenilin-1 inhibits the normal cleavage of amyloid precursor protein. *Nature* [Internet]. 1998 Jan 22;391(6665):387–90. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/9450754>
17. Vassar R, Bennett BD, Babu-Khan S, Kahn S, Mendiaz EA, Denis P, et al. Beta-secretase cleavage of Alzheimer's amyloid precursor protein by the transmembrane aspartic protease BACE. *Science* [Internet]. 1999 Oct 22 [cited 2018 Jan 18];286(5440):735–41. Available from: <http://science.sciencemag.org/content/sci/286/5440/735.full.pdf>
18. Sowade RF, Jahn TR. Seed-induced acceleration of amyloid- β mediated neurotoxicity in vivo. *Nat Commun* [Internet]. 2017 Sep 11 [cited 2017 Sep 11];8(1):512. Available from: <https://www.nature.com/articles/s41467-017-00579-4.pdf>
19. Kaye R, Head E, Thompson JL, McIntire TM, Milton SC, Cotman CW, et al. Common structure of soluble amyloid oligomers implies common mechanism of pathogenesis. *Science* [Internet]. 2003 Apr 18;300(5618):486–9. Available from:

- <http://www.ncbi.nlm.nih.gov/pubmed/12702875>
20. Heneka MT, Carson MJ, El Khoury J, Landreth GE, Brosseron F, Feinstein DL, et al. Neuroinflammation in Alzheimer's disease. *Lancet Neurol* [Internet]. 2015 Apr [cited 2018 Jan 18];14(4):388–405. Available from: https://ac.els-cdn.com/S1474442215700165/1-s2.0-S1474442215700165-main.pdf?_tid=33e16f70-fc70-11e7-892f-00000aacb362&acdnat=1516294629_3250bf4d228b6a9927dd412690cd4c73
 21. Lee VM, Goedert M, Trojanowski JQ. Neurodegenerative tauopathies. *Annu Rev Neurosci* [Internet]. 2001 Mar;24(1):1121–59. Available from: <http://www.annualreviews.org/doi/10.1146/annurev.neuro.24.1.1121>
 22. Rudrabhatla P, Jaffe H, Pant HC. Direct evidence of phosphorylated neuronal intermediate filament proteins in neurofibrillary tangles (NFTs): phosphoproteomics of Alzheimer's NFTs. *FASEB J* [Internet]. 2011 Nov;25(11):3896–905. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/21828286>
 23. Andreadis A, Brown WM, Kosik KS. Structure and novel exons of the human tau gene. *Biochemistry* [Internet]. 1992 Nov 3 [cited 2018 Jan 19];31(43):10626–33. Available from: <http://pubs.acs.org/doi/pdf/10.1021/bi00158a027>
 24. Wang Y, Mandelkow E. Tau in physiology and pathology. *Nat Rev Neurosci* [Internet]. 2016 Jan;17(1):5–21. Available from: <http://dx.doi.org/10.1038/nrn.2015.1>
 25. Rosenberg KJ, Ross JL, Feinstein HE, Feinstein SC, Israelachvili J. Complementary dimerization of microtubule-associated tau protein: Implications for microtubule bundling and tau-mediated pathogenesis. *Proc Natl Acad Sci U S A* [Internet]. 2008 May 27 [cited 2018 Jan 19];105(21):7445–50. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2396711/pdf/zpq7445.pdf>
 26. Buée L, Bussi re T, Bu e-Scherrer V, Delacourte A, Hof PR. Tau protein isoforms, phosphorylation and role in neurodegenerative disorders. *Brain Res Brain Res Rev* [Internet]. 2000 Aug;33(1):95–130. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/10967355>

27. Götz J, Ittner LM. Animal models of Alzheimer's disease and frontotemporal dementia. *Nat Rev Neurosci* [Internet]. 2008 Jul;9(7):532–44. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/18568014>
28. Hasegawa M, Morishima-Kawashima M, Takio K, Suzuki M, Titani K, Ihara Y. Protein sequence and mass spectrometric analyses of tau in the Alzheimer's disease brain. *J Biol Chem* [Internet]. 1992 Aug 25;267(24):17047–54. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/1512244>
29. Köpke E, Tung YC, Shaikh S, Alonso AC, Iqbal K, Grundke-Iqbal I. Microtubule-associated protein tau. Abnormal phosphorylation of a non-paired helical filament pool in Alzheimer disease. *J Biol Chem* [Internet]. 1993 Nov 15;268(32):24374–84. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/8226987>
30. Alonso A, Zaidi T, Novak M, Grundke-Iqbal I, Iqbal K. Hyperphosphorylation induces self-assembly of tau into tangles of paired helical filaments/straight filaments. *Proc Natl Acad Sci U S A* [Internet]. 2001 Jun 5;98(12):6923–8. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/11381127>
31. Arendt T, Stieler J, Strijkstra AM, Hut RA, Rüdiger J, Van der Zee EA, et al. Reversible paired helical filament-like phosphorylation of tau is an adaptive process associated with neuronal plasticity in hibernating animals. *J Neurosci* [Internet]. 2003 Aug 6 [cited 2018 Jan 20];23(18):6972–81. Available from: <http://www.jneurosci.org/content/jneuro/23/18/6972.full.pdf>
32. Berriman J, Serpell LC, Oberg KA, Fink AL, Goedert M, Crowther RA. Tau filaments from human brain and from in vitro assembly of recombinant protein show cross-beta structure. *Proc Natl Acad Sci U S A* [Internet]. 2003 Jul 22 [cited 2018 Jan 21];100(15):9034–8. Available from: <http://www.pnas.org/content/100/15/9034.full.pdf>
33. Fitzpatrick AWP, Falcon B, He S, Murzin AG, Murshudov G, Garringer HJ, et al. Cryo-EM structures of tau filaments from Alzheimer's disease. *Nature* [Internet]. 2017 Jul 5; Available from: <http://www.nature.com/doifinder/10.1038/nature23002>
34. Kadavath H, Hofele R V, Biernat J, Kumar S, Tepper K, Urlaub H, et al. Tau stabilizes

- microtubules by binding at the interface between tubulin heterodimers. *Proc Natl Acad Sci U S A* [Internet]. 2015 Jun 16;112(24):7501–6. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26034266><http://www.pnas.org/content/112/24/7501.full.pdf>
35. Mukrasch MD, von Bergen M, Biernat J, Fischer D, Griesinger C, Mandelkow E, et al. The “jaws” of the tau-microtubule interaction. *J Biol Chem* [Internet]. 2007 Apr 20 [cited 2018 Feb 5];282(16):12230–9. Available from: <http://www.jbc.org/content/282/16/12230.full.pdf?with-ds=yes>
 36. Wischik CM, Novak M, Thøgersen HC, Edwards PC, Runswick MJ, Jakes R, et al. Isolation of a fragment of tau derived from the core of the paired helical filament of Alzheimer disease. *Proc Natl Acad Sci U S A* [Internet]. 1988 Jun [cited 2018 Jan 21];85(12):4506–10. Available from: <http://www.pnas.org/content/85/12/4506.long>
 37. Wegmann S, Medalsy ID, Mandelkow E, Müller DJ. The fuzzy coat of pathological human Tau fibrils is a two-layered polyelectrolyte brush. *Proc Natl Acad Sci U S A* [Internet]. 2013 Jan 22;110(4):E313-21. Available from: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3557036&tool=pmcentrez&rendertype=abstract>
 38. Harada A, Oguchi K, Okabe S, Kuno J, Terada S, Ohshima T, et al. Altered microtubule organization in small-calibre axons of mice lacking tau protein. *Nature* [Internet]. 1994 Jun 9;369(6480):488–91. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/8202139>
 39. Hoover BR, Reed MN, Su J, Penrod RD, Kotilinek LA, Grant MK, et al. Tau Mislocalization to Dendritic Spines Mediates Synaptic Dysfunction Independently of Neurodegeneration. *Neuron* [Internet]. 2010;68(6):1067–81. Available from: <http://dx.doi.org/10.1016/j.neuron.2010.11.030>
 40. Goedert M, Eisenberg DS, Crowther RA. Propagation of Tau Aggregates and Neurodegeneration. *Annu Rev Neurosci* [Internet]. 2017 Jul 25 [cited 2017 Aug 23];40(1):189–210. Available from: <https://doi.org/10.1146/annurev-neuro-072116->

031153210.

41. Jack CR, Holtzman DM. Biomarker modeling of Alzheimer's disease. *Neuron* [Internet]. 2013 Dec 18 [cited 2014 Nov 7];80(6):1347–58. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24360540>
42. Iba M, McBride JD, Guo JL, Zhang B, Trojanowski JQ, Lee VM-Y. Tau pathology spread in PS19 tau transgenic mice following locus coeruleus (LC) injections of synthetic tau fibrils is determined by the LC's afferent and efferent connections. *Acta Neuropathol* [Internet]. 2015 Sep;130(3):349–62. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26150341>
43. Clavaguera F, Bolmont T, Crowther RA, Abramowski D, Frank S, Probst A, et al. Transmission and spreading of tauopathy in transgenic mouse brain. *Nat Cell Biol* [Internet]. 2009 Jul;11(7):909–13. Available from: <http://dx.doi.org/10.1038/ncb1901>
44. Clavaguera F, Akatsu H, Fraser G, Crowther RA, Frank S, Hench J, et al. Brain homogenates from human tauopathies induce tau inclusions in mouse brain. *Proc Natl Acad Sci U S A* [Internet]. 2013 Jun 4 [cited 2014 Dec 11];110(23):9535–40. Available from: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3677441&tool=pmcentrez&rendertype=abstract>
45. Iba M, Guo JL, McBride JD, Zhang B, Trojanowski JQ, Lee VM-Y. Synthetic tau fibrils mediate transmission of neurofibrillary tangles in a transgenic mouse model of Alzheimer's-like tauopathy. *J Neurosci* [Internet]. 2013 Jan 16 [cited 2015 Jan 8];33(3):1024–37. Available from: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3575082&tool=pmcentrez&rendertype=abstract>
46. Guo JL, Narasimhan S, Changolkar L, He Z, Stieber A, Zhang B, et al. Unique pathological tau conformers from Alzheimer's brains transmit tau pathology in nontransgenic mice. *J Exp Med* [Internet]. 2016 Nov 14;213(12):2635–54. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27810929>

47. Fagan AM, Mintun MA, Shah AR, Aldea P, Roe CM, Mach RH, et al. Cerebrospinal fluid tau and ptau(181) increase with cortical amyloid deposition in cognitively normal individuals: implications for future clinical trials of Alzheimer's disease. *EMBO Mol Med* [Internet]. 2009 Nov [cited 2018 Jan 25];1(8–9):371–80. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2806678/pdf/emmm0001-0371.pdf>
48. De Vos A, Bjerke M, Brouns R, De Roeck N, Jacobs D, Van den Abbeele L, et al. Neurogranin and tau in cerebrospinal fluid and plasma of patients with acute ischemic stroke. *BMC Neurol* [Internet]. 2017 Aug 30 [cited 2018 Jan 25];17(1):170. Available from: <https://bmcneurol.biomedcentral.com/track/pdf/10.1186/s12883-017-0945-8?site=bmcneurol.biomedcentral.com>
49. Galasko D, Chang L, Motter R, Clark CM, Kaye J, Knopman D, et al. High cerebrospinal fluid tau and low amyloid beta42 levels in the clinical diagnosis of Alzheimer disease and relation to apolipoprotein E genotype. *Arch Neurol* [Internet]. 1998 Jul;55(7):937–45. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/9678311>
50. Shaw LM, Vanderstichele H, Knapik-Czajka M, Clark CM, Aisen PS, Petersen RC, et al. Cerebrospinal fluid biomarker signature in Alzheimer's disease neuroimaging initiative subjects. *Ann Neurol* [Internet]. 2009 Apr;65(4):403–13. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/19296504>
51. Visser PJ, Verhey F, Knol DL, Scheltens P, Wahlund L-O, Freund-Levi Y, et al. Prevalence and prognostic value of CSF markers of Alzheimer's disease pathology in patients with subjective cognitive impairment or mild cognitive impairment in the DESCRIPA study: a prospective cohort study. *Lancet Neurol* [Internet]. 2009 Jul [cited 2018 Jan 26];8(7):619–27. Available from: https://ac.els-cdn.com/S1474442209701395/1-s2.0-S1474442209701395-main.pdf?_tid=380ecb6e-027f-11e8-9e46-00000aacb361&acdnat=1516960786_7698af7582c8285666fdbbb099b5ad2e
52. Mattsson N, Zetterberg H, Hansson O, Andreasen N, Parnetti L, Jonsson M, et al. CSF biomarkers and incipient Alzheimer disease in patients with mild cognitive

- impairment. JAMA [Internet]. 2009 Jul 22;302(4):385–93. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/19622817>
53. Mohamed NV, Herrou T, Plouffe V, Piperno N, Leclerc N. Spreading of tau pathology in Alzheimer's disease by cell-to-cell transmission. *Eur J Neurosci*. 2013;37(12):1939–48.
 54. Pooler AM, Phillips EC, Lau DHW, Noble W, Hanger DP. Physiological release of endogenous tau is stimulated by neuronal activity. *EMBO Rep* [Internet]. 2013 Apr [cited 2014 Nov 20];14(4):389–94. Available from: <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3615658&tool=pmcentrez&rendertype=abstract>
 55. Rabouille C. Pathways of Unconventional Protein Secretion. *Trends Cell Biol* [Internet]. 2017 Mar 15;27(3):230–40. Available from: <http://dx.doi.org/10.1016/j.tcb.2016.11.007>
 56. Dimou E, Nickel W. Unconventional mechanisms of eukaryotic protein secretion. *Curr Biol* [Internet]. 2018 Apr 23 [cited 2018 Apr 23];28(8):R406–10. Available from: https://ac.els-cdn.com/S0960982217315993/1-s2.0-S0960982217315993-main.pdf?_tid=47b4eb06-f68e-4465-9d7a-54afcc370c3e&acdnat=1524499162_5d0fba0aba96ec44a83be470bd5a42af
 57. Wang Y, Balaji V, Kaniyappan S, Krüger L, Irsen S, Tepper K, et al. The release and trans-synaptic transmission of Tau via exosomes. *Mol Neurodegener* [Internet]. 2017 Jan 13;12(1):5. Available from: <http://dx.doi.org/10.1186/s13024-016-0143-y>
 58. Katsinelos T, Zeitler M, Dimou E, Karakatsani A, Müller H-M, Nachman E, et al. Unconventional Secretion Mediates the Trans-cellular Spreading of Tau. *Cell Rep* [Internet]. 2018 May 15;23(7):2039–55. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/29768203>
 59. Merezko M, Brunello CA, Yan X, Vihinen H, Jokitalo E, Uronen R-L, et al. Secretion of Tau via an Unconventional Non-vesicular Mechanism. *Cell Rep* [Internet]. 2018 Nov 20;25(8):2027-2035.e4. Available from:

<https://linkinghub.elsevier.com/retrieve/pii/S2211124718316826>

60. Fulop T, Witkowski JM, Bourgade K, Khalil A, Zerif E, Larbi A, et al. Can an Infection Hypothesis Explain the Beta Amyloid Hypothesis of Alzheimer's Disease? *Front Aging Neurosci* [Internet]. 2018;10(July):224. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/30087609>
61. Eimer WA, Vijaya Kumar DK, Navalpur Shanmugam NK, Rodriguez AS, Mitchell T, Washicosky KJ, et al. Alzheimer's Disease-Associated β -Amyloid Is Rapidly Seeded by Herpesviridae to Protect against Brain Infection. *Neuron* [Internet]. 2018 Jul 11;99(1):56-63.e3. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/30001512>
62. Soscia SJ, Kirby JE, Washicosky KJ, Tucker SM, Ingelsson M, Hyman B, et al. The Alzheimer's disease-associated amyloid beta-protein is an antimicrobial peptide. *PLoS One* [Internet]. 2010 Mar 3;5(3):e9505. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/20209079>
63. Kumar DKV, Choi SH, Washicosky KJ, Eimer WA, Tucker S, Ghofrani J, et al. Amyloid- β peptide protects against microbial infection in mouse and worm models of Alzheimer's disease. *Sci Transl Med* [Internet]. 2016 May 25;8(340):340ra72. Available from: <http://stm.sciencemag.org/cgi/doi/10.1126/scitranslmed.aaf1059>
64. White MR, Kandel R, Tripathi S, Condon D, Qi L, Taubenberger J, et al. Alzheimer's associated β -amyloid protein inhibits influenza A virus and modulates viral interactions with phagocytes. *PLoS One* [Internet]. 2014;9(7):e101364. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24988208>
65. Gosztyla ML, Brothers HM, Robinson SR. Alzheimer's Amyloid- β is an Antimicrobial Peptide: A Review of the Evidence. *J Alzheimers Dis* [Internet]. 2018;62(4):1495–506. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/29504537>
66. Kountouras J, Boziki M, Polyzos SA, Katsinelos P, Gavalas E, Zeglinas C, et al. The Emerging Role of Helicobacter Pylori-Induced Metabolic Gastrointestinal Dysmotility and Neurodegeneration. *Curr Mol Med* [Internet]. 2017 Feb 7 [cited 2019 Mar 1];17(6):389–404. Available from: <http://www.eurekaselect.com/158408/article>

67. Deretzi G, Kountouras J, Polyzos SA, Zavos C, Giartza-Taxidou E, Gavalas E, et al. Gastrointestinal immune system and brain dialogue implicated in neuroinflammatory and neurodegenerative diseases. *Curr Mol Med* [Internet]. 2011 Nov;11(8):696–707. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/21902649>
68. Jiang C, Li G, Huang P, Liu Z, Zhao B. The Gut Microbiota and Alzheimer's Disease. *J Alzheimers Dis* [Internet]. 2017;58(1):1–15. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28372330>
69. Vogt NM, Romano KA, Darst BF, Engelman CD, Johnson SC, Carlsson CM, et al. The gut microbiota-derived metabolite trimethylamine N-oxide is elevated in Alzheimer's disease. *Alzheimers Res Ther* [Internet]. 2018 Dec 22;10(1):124. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/30579367>
70. Doulberis M, Kotronis G, Gialamprinou D, Kountouras J, Katsinelos P. Non-alcoholic fatty liver disease: An update with special focus on the role of gut microbiota. *Metabolism* [Internet]. 2017;71:182–97. Available from: <http://dx.doi.org/10.1016/j.metabol.2017.03.013>
71. Minter MR, Zhang C, Leone V, Ringus DL, Zhang X, Oyler-Castrillo P, et al. Antibiotic-induced perturbations in gut microbial diversity influences neuro-inflammation and amyloidosis in a murine model of Alzheimer's disease. *Sci Rep* [Internet]. 2016 Jul 21;6(June):30028. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27443609>
72. Wu S-C, Cao Z-S, Chang K-M, Juang J-L. Intestinal microbial dysbiosis aggravates the progression of Alzheimer's disease in *Drosophila*. *Nat Commun* [Internet]. 2017;8(1):24. Available from: <http://dx.doi.org/10.1038/s41467-017-00040-6>
73. Clemente JC, Ursell LK, Parfrey LW, Knight R. The impact of the gut microbiota on human health: an integrative view. *Cell* [Internet]. 2012 Mar 16;148(6):1258–70. Available from: <http://dx.doi.org/10.1016/j.cell.2012.01.035>
74. Alkasir R, Li J, Li X, Jin M, Zhu B. Human gut microbiota: the links with dementia development. *Protein Cell* [Internet]. 2017;8(2):90–102. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27866330>

75. Friedland RP, Chapman MR. The role of microbial amyloid in neurodegeneration. *PLoS Pathog* [Internet]. 2017;13(12):e1006654. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/29267402>
76. Zhuang Z-Q, Shen L-L, Li W-W, Fu X, Zeng F, Gui L, et al. Gut Microbiota is Altered in Patients with Alzheimer's Disease. *J Alzheimers Dis* [Internet]. 2018;63(4):1337–46. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/29758946>
77. Saji N, Niida S, Murotani K, Hisada T, Tsuduki T, Sugimoto T, et al. Analysis of the relationship between the gut microbiome and dementia: a cross-sectional study conducted in Japan. *Sci Rep* [Internet]. 2019 Jan 30;9(1):1008. Available from: <http://dx.doi.org/10.1038/s41598-018-38218-7>
78. Vogt NM, Kerby RL, Dill-McFarland KA, Harding SJ, Merluzzi AP, Johnson SC, et al. Gut microbiome alterations in Alzheimer's disease. *Sci Rep* [Internet]. 2017;7(1):13537. Available from: <http://dx.doi.org/10.1038/s41598-017-13601-y>
79. Cattaneo A, Cattane N, Galluzzi S, Provasi S, Lopizzo N, Festari C, et al. Association of brain amyloidosis with pro-inflammatory gut bacterial taxa and peripheral inflammation markers in cognitively impaired elderly. *Neurobiol Aging* [Internet]. 2017;49:60–8. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27776263>
80. MahmoudianDehkordi S, Arnold M, Nho K, Ahmad S, Jia W, Xie G, et al. Altered bile acid profile associates with cognitive impairment in Alzheimer's disease-An emerging role for gut microbiome. *Alzheimers Dement* [Internet]. 2019;15(1):76–92. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/30337151>
81. Lee Y-L, Hu H-Y, Huang L-Y, Chou P, Chu D. Periodontal Disease Associated with Higher Risk of Dementia: Population-Based Cohort Study in Taiwan. *J Am Geriatr Soc* [Internet]. 2017 Sep;65(9):1975–80. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/28598507>
82. Hayashi K, Hasegawa Y, Takemoto Y, Cao C, Takeya H, Komohara Y, et al. Continuous intracerebroventricular injection of *Porphyromonas gingivalis* lipopolysaccharide induces systemic organ dysfunction in a mouse model of

- Alzheimer's disease. *Exp Gerontol* [Internet]. 2019 Jun;120(February):1–5. Available from: <https://doi.org/10.1016/j.exger.2019.02.007>
83. Choi S, Kim K, Chang J, Kim SM, Kim SJ, Cho H-J, et al. Association of Chronic Periodontitis on Alzheimer's Disease or Vascular Dementia. *J Am Geriatr Soc* [Internet]. 2019 Jun;67(6):1234–9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/30874308>
 84. Roubaud-Baudron C, Krolak-Salmon P, Quadrio I, Mégraud F, Salles N. Impact of chronic *Helicobacter pylori* infection on Alzheimer's disease: preliminary results. *Neurobiol Aging* [Internet]. 2012 May;33(5):1009.e11-9. Available from: <http://dx.doi.org/10.1016/j.neurobiolaging.2011.10.021>
 85. Beydoun MA, Beydoun HA, Shroff MR, Kitner-Triolo MH, Zonderman AB. *Helicobacter pylori* seropositivity and cognitive performance among US adults: evidence from a large national survey. *Psychosom Med* [Internet]. 2013 Jun [cited 2019 Aug 7];75(5):486–96. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/23697465>
 86. Boziki M, Polyzos SA, Deretzi G, Kazakos E, Katsinelos P, Doulberis M, et al. A potential impact of *Helicobacter pylori*-related galectin-3 in neurodegeneration. *Neurochem Int* [Internet]. 2018;113:137–51. Available from: <https://doi.org/10.1016/j.neuint.2017.12.003>
 87. Rea K, Dinan TG, Cryan JF. The microbiome: A key regulator of stress and neuroinflammation. *Neurobiol Stress* [Internet]. 2016 Oct;4:23–33. Available from: <http://dx.doi.org/10.1016/j.ynstr.2016.03.001>
 88. Zárate-Bladés CR, Horai R, Caspi RR. Regulation of Autoimmunity by the Microbiome. *DNA Cell Biol* [Internet]. 2016 Sep;35(9):455–8. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27463238>
 89. Stanisavljević S, Lukić J, Soković S, Mihajlovic S, Mostarica Stojković M, Miljković D, et al. Correlation of Gut Microbiota Composition with Resistance to Experimental Autoimmune Encephalomyelitis in Rats. *Front Microbiol* [Internet]. 2016;7(December):2005. Available from:

<http://www.ncbi.nlm.nih.gov/pubmed/28018327>

90. Tükel C, Wilson RP, Nishimori JH, Pezeshki M, Chromy BA, Bäumler AJ. Responses to amyloids of microbial and host origin are mediated through toll-like receptor 2. *Cell Host Microbe* [Internet]. 2009 Jul 23;6(1):45–53. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/19616765>
91. Holmes C. Review: systemic inflammation and Alzheimer's disease. *Neuropathol Appl Neurobiol* [Internet]. 2013 Feb;39(1):51–68. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/23046210>
92. César Machado MC, da Silva FP. Intestinal Barrier Dysfunction in Human Pathology and Aging. *Curr Pharm Des* [Internet]. 2016 Sep 7 [cited 2019 Aug 7];22(30):4645–50. Available from: <http://www.eurekaselect.com/openurl/content.php?genre=article&issn=1381-6128&volume=22&issue=30&spage=4645>
93. Jakobsson HE, Rodríguez-Piñeiro AM, Schütte A, Ermund A, Boysen P, Bemark M, et al. The composition of the gut microbiota shapes the colon mucus barrier. *EMBO Rep* [Internet]. 2015 Feb;16(2):164–77. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25525071>
94. Kowalski K, Mulak A. Brain-Gut-Microbiota Axis in Alzheimer's Disease. *J Neurogastroenterol Motil* [Internet]. 2019 Jan 31;25(1):48–60. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/30646475>
95. Bostanciklioğlu M. The role of gut microbiota in pathogenesis of Alzheimer's disease. *J Appl Microbiol* [Internet]. 2019 Mar 28;1–14. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/30920075>
96. Chen R, Wu P, Cai Z, Fang Y, Zhou H, Lasanajak Y, et al. Puerariae Lobatae Radix with Chuanxiong Rhizoma for treatment of cerebral ischemic stroke by remodeling gut microbiota to regulate the brain-gut barriers of dietary capsaicin against chronic low-grade inflammation. *J Nutr Biochem* [Internet]. 2018 Dec 21;65:101–14. Available from: <https://doi.org/10.1016/j.jnutbio.2018.12.004>

97. Marungruang N, Arévalo Sureda E, Lefrançoise A, Weström B, Nyman M, Prykhodko O, et al. Impact of dietary induced precocious gut maturation on cecal microbiota and its relation to the blood-brain barrier during the postnatal period in rats. *Neurogastroenterol Motil* [Internet]. 2018 Jun;30(6):e13285. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/29327435>
98. Roy Sarkar S, Banerjee S. Gut microbiota in neurodegenerative disorders. *J Neuroimmunol* [Internet]. 2019 Mar 15;328(December 2018):98–104. Available from: <https://doi.org/10.1016/j.jneuroim.2019.01.004>
99. Cheng Y, Bai F. The Association of Tau With Mitochondrial Dysfunction in Alzheimer's Disease. *Front Neurosci* [Internet]. 2018;12(MAR):163. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/29623026>
100. Syeda T, Sanchez-Tapia M, Pinedo-Vargas L, Granados O, Cuervo-Zanatta D, Rojas-Santiago E, et al. Bioactive Food Abates Metabolic and Synaptic Alterations by Modulation of Gut Microbiota in a Mouse Model of Alzheimer's Disease. *J Alzheimers Dis* [Internet]. 2018;66(4):1657–82. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/30475761>
101. Li J, Perez-Perez GI. *Helicobacter pylori* the Latent Human Pathogen or an Ancestral Commensal Organism. *Front Microbiol* [Internet]. 2018;9(APR):609. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/29666614>
102. Franceschi F, Ojetti V, Candelli M, Covino M, Cardone S, Potenza A, et al. Microbes and Alzheimer' disease: lessons from *H. pylori* and GUT microbiota. *Eur Rev Med Pharmacol Sci* [Internet]. 2019 Jan;23(1):426–30. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/30657587>
103. He C, Yang Z, Lu N. Imbalance of Gastrointestinal Microbiota in the Pathogenesis of *Helicobacter pylori*-Associated Diseases. *Helicobacter* [Internet]. 2016 Oct;21(5):337–48. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26876927>
104. Bruno G, Rocco G, Zaccari P, Porowska B, Mascellino MT, Severi C. *Helicobacter pylori* Infection and Gastric Dysbiosis: Can Probiotics Administration Be Useful to

- Treat This Condition? *Can J Infect Dis Med Microbiol = J Can des Mal Infect la Microbiol medicale* [Internet]. 2018;2018:6237239. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/30275917>
105. Shen L, Liu L, Ji H-F. Alzheimer's Disease Histological and Behavioral Manifestations in Transgenic Mice Correlate with Specific Gut Microbiome State. *J Alzheimers Dis* [Internet]. 2017;56(1):385–90. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/27911317>
 106. Katan M, Moon YP, Paik MC, Sacco RL, Wright CB, Elkind MS V. Infectious burden and cognitive function: the Northern Manhattan Study. *Neurology* [Internet]. 2013 Mar 26;80(13):1209–15. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/23530151>
 107. Wright CB, Gardener H, Dong C, Yoshita M, DeCarli C, Sacco RL, et al. Infectious Burden and Cognitive Decline in the Northern Manhattan Study. *J Am Geriatr Soc* [Internet]. 2015 Aug;63(8):1540–5. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/26289683>
 108. Xu Y, Wang Q, Liu Y, Cui R, Lu K, Zhao Y. Association between *Helicobacter pylori* infection and carotid atherosclerosis in patients with vascular dementia. *J Neurol Sci* [Internet]. 2016 Mar 15;362:73–7. Available from: <http://dx.doi.org/10.1016/j.jns.2016.01.025>
 109. Swardfager W, Lanctôt K, Rothenburg L, Wong A, Cappell J, Herrmann N. A meta-analysis of cytokines in Alzheimer's disease. *Biol Psychiatry* [Internet]. 2010 Nov 15;68(10):930–41. Available from: <http://dx.doi.org/10.1016/j.biopsych.2010.06.012>
 110. Beydoun MA, Beydoun HA, Elbejjani M, Dore GA, Zonderman AB. *Helicobacter pylori* seropositivity and its association with incident all-cause and Alzheimer's disease dementia in large national surveys. *Alzheimers Dement* [Internet]. 2018 Sep;14(9):1148–58. Available from: <https://doi.org/10.1016/j.jalz.2018.04.009>
 111. Kountouras J, Mylopoulos N, Chatzopoulos D, Zavos C, Boura P, Konstas AGP, et al. Eradication of *Helicobacter pylori* may be beneficial in the management of chronic open-angle glaucoma. *Arch Intern Med* [Internet]. 2002 Jun 10;162(11):1237–44.

Available from: <http://www.ncbi.nlm.nih.gov/pubmed/12038941>

112. Kountouras J, Zavos C, Gavalas E, Boziki M, Chatzopoulos D, Katsinelos P. Normal-tension glaucoma and Alzheimer's disease: *Helicobacter pylori* as a possible common underlying risk factor. *Med Hypotheses* [Internet]. 2007;68(1):228–9. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/16978795>
113. Supajatura V, Ushio H, Wada A, Yahiro K, Okumura K, Ogawa H, et al. Cutting edge: VacA, a vacuolating cytotoxin of *Helicobacter pylori*, directly activates mast cells for migration and production of proinflammatory cytokines. *J Immunol* [Internet]. 2002 Mar 15;168(6):2603–7. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/11884423>
114. McKittrick CM, Lawrence CE, Carswell HVO. Mast cells promote blood brain barrier breakdown and neutrophil infiltration in a mouse model of focal cerebral ischemia. *J Cereb Blood Flow Metab* [Internet]. 2015 Mar 31;35(4):638–47. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25564235>
115. Wang X-L, Zeng J, Feng J, Tian Y-T, Liu Y-J, Qiu M, et al. *Helicobacter pylori* filtrate impairs spatial learning and memory in rats and increases β -amyloid by enhancing expression of presenilin-2. *Front Aging Neurosci* [Internet]. 2014;6(April):66. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/24782763>
116. Wang X-L, Zeng J, Yang Y, Xiong Y, Zhang Z-H, Qiu M, et al. *Helicobacter pylori* filtrate induces Alzheimer-like tau hyperphosphorylation by activating glycogen synthase kinase-3 β . *J Alzheimers Dis* [Internet]. 2015;43(1):153–65. Available from: <http://www.ncbi.nlm.nih.gov/pubmed/25079798>